

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number
WO 01/04139 A2

(51) International Patent Classification?: C07K
(21) International Application Number: PCT/US00/19001
(22) International Filing Date: 13 July 2000 (13.07.2000)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
9916417.0 13 July 1999 (13.07.1999) GB
60/169,573 8 December 1999 (08.12.1999) US

(71) Applicants (for all designated States except US):
SMITHKLINE BEECHAM CORPORATION
[US/US]; One Franklin Plaza, Philadelphia, PA 19103
(US). SMITHKLINE BEECHAM PLC [GB/GB]; New
Horizons Court, Great West Road, Brentford, Middlesex
TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): AMES, Robert, S.
[US/US]; 276 Ellis Road, Havertown, PA 19083 (US).

ELSHOURBAGY, Nabil [US/US]; 1648 Bow Tree
Drive, West Chester, PA 19380 (US). FOLEY, James, J.
[US/US]; 117 Timothy Circle, Radnor, PA 19087 (US).
MICHALOVICH, David [GB/GB]; 89c Ferme Park
Road, Crouch End, London N8 9SA (GB). SARAU,
Henry, M. [US/US]; 348 Maple Avenue, Harleysville,
PA 19438 (US). SMITH, Randall [US/US]; 4138 Pres-
idential Drive, Lafayette Hill, PA 19444 (US). TSUI,
Ping [US/US]; 1237 Berwyn-Paoli Road, Berwyn, PA
19312 (US). VAWTER, Lisa [US/US]; 1075 Ebert Road,
Coopersburg, PA 18036 (US). AGARWAL, Pankaj
[IN/US]; 251 W. DeKalb Pike, E801, King of Prussia, PA
19406 (US). LANE, Pamela [US/US]; 31 Colton Drive,
Norristown, PA 19401 (US).

(74) Agents: ANDERSEN, Robert, L. et al.; Ratner & Prestia,
301 One Westlakes (Berwyn), P.O. Box 980, Valley Forge,
PA 19482-0980 (US).

(81) Designated States (national): JP, US.

(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

[Continued on next page]

(54) Title: POLYNUCLEOTIDE AND POLYPEPTIDE SEQUENCES OF HUMAN AXOR29 RECEPTOR AND METHODS OF
SCREENING FOR AGONISTS AND ANTAGONISTS OF THE INTERACTION BETWEEN HUMAN AXOR29 RECEPTOR
AND ITS LIGANDS

ATGGAGTCGGGCTGCTGCGCGCGCGCGGTGAGCGAGGTGATCGTCTGCATTACAAC
TACACCGCAAGCTCCGCGGTGCGCTACACGCGCGGTGCGCGCTGCGCGCGCGCGC
GTGGTGTGCTGCGGTGCGGTGCGGTGCGGTGCGGTGCGGTGCGGTGCGGTGCGGTG
CTCGGACGCCACCGCGCTTCCACGCTCCCATGTTCTGCTCTGCGGCGAGCTCACGTTG
TCGGATCTGCTGCGGAGCGCGCTACGCGCGCAACATCTACTGTGCGGCGCGCTCACG
CTGAACCTGTCCCGCGCTCTGGTTGCGACGCGGAGGAGCGCTCTGCTGGCACTCACT
CGCTCGGTGCTGAGCCTCTGCGCATGCGCTGAGCGCGCTCACCATGCGCGCGAGG
GGGCGCGCGCGCTCTCAGTCCGCGCGCGCGCTGCGGATGCGAGCGCGCGCTGCGG
GTGTGCTGCTCTCGGCTCTGCGCGCGCTGCGCGCGCTGCGCGCGCTGCGCGCGCTG
GCTTGTCTCACTGTCTTGGCGCTCTAGCGCAAGGCTACGTGCTCTTCTGCTGCTCGCC
TTCGTGGGATCTGCGCGCTATCTGTGCACTCTACGCGCGCTCTACTGCCAGGTACGC
GCCAAGCGCGCGCTGCGCGCACGCGCGCGGACTGCGGAGCACCTCGACCGCGCGG
CGTCGCAAGCGCGCTGCTGCGCTTGTGCGCGCTGCGCGCGCTGCGCGCGCTGCGCG
GTGGCATGTTGGGCGCGCTCTTCTGCTGCTGTTGCTGCGAGCTGCGCGTGGCGCGCG
ACCTGTCTGTACTCTGCGAGCGGATCCCTTCTGCGGACTGCGCATGCGCAACTCACTT
CTGAACCCCATCATCTACAGCTCACCACCGCGACTGCGCGCGCGCTCTGCGCGCTG
GTCTGCTGCGGACGCCACTCTGCGCGAGAGACCGAGTGGCTCCAGCACTGCGCGAGC
GCGGCTGAGGCTTCCGCGCGCTGCGCGCGCTGCGCGCGCTGCGCGCGCTGCGCGCGCTT
AGCGGCTGCGGAGCGCTCTGCGCGCGCGCGCTGCGCGCGCTGCGCGCGCTTCCAGGCG
AGCCCGCGTGCACCCACAGCGCGCGCGCTGCGTGTATCAGAACCGCGCTGCGAGTGA

(57) Abstract: Human AXOR29 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Human AXOR29 is identified as a selective receptor for sphingosine-1-phosphate ("S-1-P") and di-hydro sphingosine-1-phosphate ("di-hydro S-1-P"). Also disclosed are methods for discovering agonists and antagonists of the interaction between S-1-P and di-hydro S-1-P and their cellular receptor, human AXOR29, which may have utility in the treatment of several human diseases and disorders, including, but not limited to the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; stroke; ulcers; asthma; allergic; benign prostatic hypertrophy; migraine; vomiting; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, depression, delirium, dementia, and severe mental retardation.



Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

- 1 -

**POLYNUCLEOTIDE AND POLYPEPTIDE SEQUENCES OF HUMAN AXOR29
RECEPTOR AND METHODS OF SCREENING FOR AGONISTS AND
ANTAGONISTS OF THE INTERACTION BETWEEN HUMAN
AXOR29 RECEPTOR AND ITS LIGANDS**

5 FIELD OF THE INVENTION

This invention relates to newly identified polypeptides and polynucleotides encoded by them and to the use of such polynucleotides and polypeptides, and to their production. More particularly, the polynucleotides and polypeptides of the present invention relate to the G-protein coupled receptors, hereinafter referred to as human AXOR29 receptor. This invention also relates to
10 methods for discovering agonists and antagonists of the interaction between sphingosine 1-phosphate (hereinafter referred to as "S-1-P") and di-hydro sphingosine-1-phosphate (hereinafter referred to as "di-hydro S-1-P") and their cellular receptor, human AXOR29 receptor. The invention also relates to the use of human AXOR29 receptor polynucleotides and polypeptides in therapy and in identifying compounds which may be agonists, antagonists and /or inhibitors which
15 are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

BACKGROUND OF THE INVENTION

The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superseding earlier
20 approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular
25 biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

It is well established that many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers, e.g., cAMP (Lefkowitz, *Nature*, 1991, 351:353-354). Herein these proteins are referred to
30 as proteins participating in pathways with G-proteins or PPG proteins. Some examples of these proteins include the GPC receptors, such as those for adrenergic agents and dopamine (Kobilka, B.K., *et al.*, *Proc. Natl Acad. Sci., USA*, 1987, 84:46-50; Kobilka, B.K., *et al.*, *Science*, 1987, 238:650-656; Bunzow, J.R., *et al.*, *Nature*, 1988, 336:783-787), G-proteins themselves, effector proteins, e.g.,

- 2 -

phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins, *e.g.*, protein kinase A and protein kinase C (Simon, M.I., *et al.*, *Science*, 1991, 252:802-8).

For example, in one form of signal transduction, the effect of hormone binding is activation of the enzyme, adenylyl cyclase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide GTP. GTP also influences hormone binding. A G-protein connects the hormone receptor to adenylyl cyclase. G-protein was shown to exchange GTP for bound GDP when activated by a hormone receptor. The GTP-carrying form then binds to activated adenylyl cyclase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

The membrane protein gene superfamily of G-protein coupled receptors has been characterized as having seven putative transmembrane domains. The domains are believed to represent transmembrane α -helices connected by extracellular or cytoplasmic loops. G-protein coupled receptors include a wide range of biologically active receptors, such as hormone, viral, growth factor and neuroreceptors.

G-protein coupled receptors (otherwise known as 7TM receptors) have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. The G-protein family of coupled receptors includes dopamine receptors which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include, but are not limited to, calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorant, and cytomegalovirus receptors.

Most G-protein coupled receptors have single conserved cysteine residues in each of the first two extracellular loops which form disulfide bonds that are believed to stabilize functional protein structure. The 7 transmembrane regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 has been implicated in signal transduction.

Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues can influence signal transduction of some G-protein coupled receptors. Most G-protein coupled receptors contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus. For several G-protein coupled receptors, such as the β -adrenoreceptor, phosphorylation by protein kinase A and/or specific receptor kinases mediates receptor desensitization.

- 3 -

For some receptors, the ligand binding sites of G-protein coupled receptors are believed to comprise hydrophilic sockets formed by several G-protein coupled receptor transmembrane domains, said socket being surrounded by hydrophobic residues of the G-protein coupled receptors. The hydrophilic side of each G-protein coupled receptor transmembrane helix is postulated to face inward and form polar ligand binding site. TM3 has been implicated in several G-protein coupled receptors as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine and TM6 or TM7 phenylalanines or tyrosines are also implicated in ligand binding.

G-protein coupled receptors can be intracellularly coupled by heterotrimeric G-proteins to various intracellular enzymes, ion channels and transporters (see, Johnson, *et al.*, *Endoc. Rev.*, 1989, 10:317-331). Different G-protein α -subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of G-protein coupled receptors have been identified as an important mechanism for the regulation of G-protein coupling of some G-protein coupled receptors. G-protein coupled receptors are found in numerous sites within a mammalian host.

Over the past 15 years, nearly 350 therapeutic agents targeting 7 transmembrane (7 TM) receptors have been successfully introduced onto the market.

SUMMARY OF THE INVENTION

The present invention relates to human AXOR29 receptor, in particular human AXOR29 receptor polypeptides and human AXOR29 receptor polynucleotides, recombinant materials and methods for their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain diseases, including, but not limited to infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; stroke; ulcers; asthma; allergies; benign prostatic hypertrophy; migraine; vomiting; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, depression, delirium, dementia, and severe mental retardation, hereinafter referred to as "diseases of the invention".

In accordance with another aspect of the present invention there are provided methods of screening for compounds which bind to and activate (agonist) or inhibit activation (antagonist) of human AXOR29 receptor polypeptides, and for their ligands.

In particular, the preferred method for identifying agonist or antagonist of a human AXOR29 receptor polypeptide comprises:

- 4 -

(a) contacting a cell expressing on the surface thereof the polypeptide, said polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and

5 (b) determining whether the compound binds to and activates or inhibits the polypeptide by measuring the level of a signal generated from the interaction of the compound with the polypeptide.

In a further preferred embodiment, the method further comprises conducting the identification of agonist or antagonist in the presence of labeled or unlabeled S-1-P or di-hydro S-1-P.

10 In another embodiment, the method for identifying agonist or antagonist of a human AXOR29 receptor polypeptide comprises:

determining the inhibition of binding of a ligand to cells which have the polypeptide on the surface thereof, or to cell membranes containing the polypeptide, in the presence of a candidate compound under conditions to permit binding to the polypeptide, and determining the amount of ligand bound to the polypeptide, such that a compound capable of causing reduction of binding of a ligand is an agonist or
15 antagonist. Preferably, the ligand is S-1-P or di-hydro S-1-P. Yet more preferably, S-1-P or di-hydro S-1-P is labeled.

Furthermore, the present invention relates to treating conditions associated with human AXOR29 receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate human AXOR29 receptor activity or levels.

20 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of the human AXOR29 receptor (SEQ ID NO:1).

Figure 2 shows the deduced amino acid sequence of the human AXOR29 receptor (SEQ ID NO:2).

25 Figure 3 shows concentration-response curves for S-1-P with 3 clones of the human AXOR29 receptor stably expressed in RBL 2H3 cells and also for RBL 2H3 cells and RBL 2H3 vector control cells.

Figure 4 shows concentration-response curves for lysophosphatidic acid (LPA) with 3 clones of the AXOR29 receptor stably expressed in RBL 2H3 cells and also for RBL 2H3 cells and RBL 2H3 vector control cells.

- 5 -

DESCRIPTION OF THE INVENTION

In a first aspect, the present invention relates to AXOR29 polypeptides. Such polypeptides include:

- (a) an isolated polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;
- 5 (b) an isolated polypeptide comprising a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (c) an isolated polypeptide comprising the polypeptide sequence of SEQ ID NO:2;
- (d) an isolated polypeptide having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- 10 (e) the polypeptide sequence of SEQ ID NO:2; and
- (f) an isolated polypeptide having or comprising a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2;
- (g) fragments and variants of such polypeptides in (a) to (f).

Polypeptides of the present invention are believed to be members of the 7 transmembrane G-protein coupled receptor family of polypeptides. The biological properties of the human AXOR29 receptor are hereinafter referred to as "biological activity of human AXOR29 receptor" or "human AXOR29 receptor activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of human AXOR29 receptor.

Polypeptides of the present invention also include variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

Preferred fragments of polypeptides of the present invention include an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2, or an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2. Preferred fragments are biologically active fragments that mediate the biological activity of human AXOR29 receptor, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

- 6 -

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods.. Means for preparing such polypeptides are well understood in the art.

In a further aspect, the present invention relates to human AXOR29 receptor polynucleotides. Such polynucleotides include:

- (a) an isolated polynucleotide comprising a polynucleotide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide sequence of SEQ ID NO:1;
- (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1;
- (c) an isolated polynucleotide having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide of SEQ ID NO:1;
- (d) the isolated polynucleotide of SEQ ID NO:1;
- (e) an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (f) an isolated polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2;
- (g) an isolated polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (h) an isolated polynucleotide encoding the polypeptide of SEQ ID NO:2;
- (i) an isolated polynucleotide having or comprising a polynucleotide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polynucleotide sequence of SEQ ID NO:1;

- 7 -

(j) an isolated polynucleotide having or comprising a polynucleotide sequence encoding a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2; and

polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.

Preferred fragments of polynucleotides of the present invention include an isolated polynucleotide comprising an nucleotide sequence having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1, or an isolated polynucleotide comprising an sequence having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ ID NO: 1.

Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including polynucleotides having one or more single nucleotide polymorphisms (SNPs).

Polynucleotides of the present invention also include polynucleotides encoding polypeptide variants that comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acid residues are substituted, deleted or added, in any combination.

In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide that:

(a) comprises an RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;

(b) is the RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;

(c) comprises an RNA transcript of the DNA sequence of SEQ ID NO:1; or

(d) is the RNA transcript of the DNA sequence of SEQ ID NO:1;

and RNA polynucleotides that are complementary thereto.

The polynucleotide sequence of SEQ ID NO:1 shows homology with Mouse edg-1 G-protein coupled receptor (C.H. Liu and T. Hla, Genomics 43:15-24, 1997). The polynucleotide sequence of SEQ ID NO:1 is a cDNA sequence that encodes the polypeptide of SEQ ID NO:2. The polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence of SEQ ID NO:1 or it may be a sequence other than SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

The polypeptide of the SEQ ID NO:2 is related to other proteins of the 7 transmembrane G-protein coupled receptor family, having homology and/or structural similarity with Mouse edg-1 G-protein coupled receptor (C.H. Liu and T. Hla, *Genomics* 43:15-24, 1997).

Through phylogenetic analysis, AXOR29 was identified to be an EDG receptor. EDG receptors comprise a small, closely-related group of GPCRs. Known EDG receptors are EDG-1, EDG-2, EDG-3, EDG-4, EDG-5, EDG-6 and EDG-7. See Moolenaar, *et al.*, *J. Curr. Opin. Cell Biol.* 9(2): 168-173 (1997); MacLennan, *et al.*, *J. Cell. Neurosci.* 5(3):201-209 (1994); Graler, *et al.*, *J. Genomics* 53(2): 164-169 (1998); Bandoh, *Acid. J. Biol. Chem.* 274: 27776-27785 (1999). Reports on the ligand pairing of the EDG receptors indicate that the EDG-1, -3 and -5 receptors respond to S-1-P, whereas the EDG-2, -4 and -7 receptors use LPA as their ligand. See Lee, *et al.*, *Science* 279: 1552-1555 (1998); Moolenaar, *et al.*, *supra*; Bandoh, *et al.*, *supra*; Okamoto, *et al.*, *Biotech. Biophys. Res. Comm.* 260:203 (1999); Ancellin, *et al.*, *J Bio Chem* 274: 18997 (1999). The ligand for the EDG-6 receptor has yet to be identified.

We have found the ligand usage pattern of the EDG receptors to reflect the phylogenetic relationships among these genes, identified by shared, derived mutations among the ligand-usage groups. These relationships are evidenced by the presence of shared, derived fixed mutations that are individual to each of the groups. Swofford, *et al.*, (1996) Chap. 11, PHYLOGENETIC INFERENCE IN MOLECULAR SYSTEMATICS (Eds. Hillis, D.M., C. Moritz, B.K. Mable, Sinauer Assoc., Sunderland Mass.), 655. In short, EDG-1, -3 and -5 receptors are closest relatives. Likewise, EDG-2, -4 and -7 are more closely-related to each other than any of them is to the S-1-P -responding EDG receptor group.

The human AXOR29 receptor shares 42% amino acid identity with EDG-1 and shares the same derived, fixed mutations that we identified in the S-1-P-responding group of EDG receptors (EDG-1, -3 and -5). We we have confirmed through experimentation that S-1-P is the ligand for AXOR29. See Example 9 and Figures 3 and 4.

Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one human AXOR29 receptor activity.

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human lymph, prostate, pancreas and multiple sclerosis lesions, (see for instance, Sambrook, *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

(1989)). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence, or other fusion peptide portions. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a polynucleotide sequence of SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO:1, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes isolated polynucleotides, preferably with a nucleotide sequence of at

- 10 -

least 100, obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof, preferably of at least 15 nucleotides.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *Proc Nat Acad Sci USA* 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trade mark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark) technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis, *et al.*, BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook, *et al.* (*ibid*). Preferred

- 11 -

methods of introducing polynucleotides into host cells include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression.

Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or purification.

- 12 -

Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled human AXOR29 receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton, *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401).

An array of oligonucleotides probes comprising human AXOR29 receptor polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability, see, for example, M.Chee, *et al.*, Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in

- 13 -

the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit comprising:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment or an RNA transcript thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

The polynucleotide sequences of the present invention are valuable for chromosome localisation studies. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (co-inheritance of physically adjacent genes). Precise human chromosomal localisations for a genomic sequence (gene fragment etc.) can be determined using Radiation Hybrid (RH) Mapping (Walter, M. Spillett, D., Thomas, P., Weissenbach, J., and Goodfellow, P., (1994) A method for constructing radiation hybrid maps of whole genomes, *Nature Genetics* 7, 22-28). A number of RH panels are available from Research Genetics (Huntsville, AL, USA) e.g. the GeneBridge4 RH panel (*Hum Mol Genet* 1996 Mar;5(3):339-46 A radiation hybrid map of the human genome. Gyapay G, Schmitt K, Fizames C, Jones H, Vega-Czarny N, Spillett D, Muselet D, Prud'Homme JF, Dib C, Auffray C, Morissette J, Weissenbach J, Goodfellow PN). To determine the chromosomal location of a gene using this panel, 93 PCRs are performed using primers designed from the gene of interest on RH DNAs. Each of these DNAs contains random human genomic fragments maintained in a hamster background (human / hamster hybrid cell lines). These PCRs result in 93 scores indicating the

presence or absence of the PCR product of the gene of interest. These scores are compared with scores created using PCR products from genomic sequences of known location. This comparison is conducted at <http://www.genome.wi.mit.edu/>. The gene of the present invention maps to human chromosome 19p13.2.

5 The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridisation techniques to clones arrayed on a grid, such as cDNA
10 microarray hybridisation (Schena *et al*, Science, 270, 467-470, 1995 and Shalon *et al*, Genome Res, 6, 639-645, 1996) and nucleotide amplification techniques such as PCR. A preferred method uses the TAQMAN (Trade mark) technology available from Perkin Elmer. Results from these studies can provide an indication of the normal function of the polypeptide in the organism. In addition,
15 comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by an alternative form of the same gene (for example, one having an alteration in polypeptide coding potential or a regulatory mutation) can provide valuable insights into the role of the polypeptides of the present invention, or that of inappropriate expression thereof in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

20 A further aspect of the present invention relates to antibodies. The polypeptides of the invention or their fragments, or cells expressing them, can be used as immunogens to produce antibodies that are immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

25 Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al*, *Immunology Today* (1983) 4:72) and the EBV-
30 hybridoma technique (Cole, *et al*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention.

- 15 -

Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease, whether that disease is already established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

The human AXOR29 receptor polypeptide of the present invention may be employed in a process for screening for compounds that bind to and activate the human AXOR29 receptor polypeptides of the present invention (called agonists), or inhibit the interaction of the human AXOR29 receptor polypeptides with receptor ligands (called antagonists). Thus, polypeptides of the

- 16 -

invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan, *et al.*, CURRENT PROTOCOLS IN IMMUNOLOGY 1(2):Chapter 5 (1991).

Human AXOR29 receptor proteins are responsible for many biological functions, including many pathologies. Provided by the invention are screening methods to identify compounds and drugs that stimulate human AXOR29 receptor or that inhibit the function or level of the polypeptide. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions as infections such as diseases of the invention, among others. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions as diseases of the invention.

In general, such screening procedures involve providing appropriate cells that express the receptor polypeptide of the present invention on the surface thereof. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. In particular, a polynucleotide encoding the receptor of the present invention is employed to transfect cells to thereby express the human AXOR29 receptor polypeptide. The expressed receptor is then contacted with a test compound to observe binding, stimulation or inhibition of a functional response.

One such screening procedure involves the use of melanophores that are transfected to express the human AXOR29 receptor polypeptide of the present invention. Such a screening technique is described in PCT WO 92/01810, published February 6, 1992. Such an assay may be employed to screen for a compound which inhibits activation of the receptor polypeptide of the present invention by contacting the melanophore cells that encode the receptor with both a receptor ligand, such as S-1-P, and a compound to be screened. Inhibition of the signal generated by the ligand indicates that a compound is a potential antagonist for the receptor, *i.e.*, inhibits activation of the receptor.

The technique may also be employed for screening of compounds that activate the receptor by contacting such cells with compounds to be screened and determining whether such compound generates a signal, *i.e.*, activates the receptor.

Other screening techniques include the use of cells which express the human AXOR29 receptor polypeptide (for example, transfected CHO cells) in a system that measures extracellular pH changes caused by receptor activation. In this technique, compounds may be contacted with cells expressing the receptor polypeptide of the present invention. A second messenger response, *e.g.*, signal transduction or pH changes, is then measured to determine whether the potential compound activates or inhibits the receptor.

- 17 -

Another screening technique involves expressing the human AXOR29 receptor polypeptide in which the receptor is linked to phospholipase C or D. Representative examples of such cells include, but are not limited to: endothelial cells, smooth muscle cells, and embryonic kidney cells. The screening may be accomplished as hereinabove described by detecting activation of the receptor or inhibition of activation of the receptor from the phospholipase second signal.

Another method involves screening for compounds that are antagonists, and thus inhibit activation of the receptor polypeptide of the present invention by determining inhibition of binding of labeled ligand, such as S-1-P or di-hydro S-1-P, to cells expressing the receptor on their surface, or cell membranes containing the receptor. Such a method involves transfecting a eukaryotic cell with DNA encoding the human AXOR29 receptor polypeptide, such that the cell expresses the receptor on its surface. The cell is then contacted with a potential antagonist in the presence of a labeled form of a ligand, such as S-1-P or di-hydro S-1-P. The ligand can be labeled, *e.g.*, by radioactivity. The amount of labeled ligand bound to the receptors is measured, *e.g.*, by measuring radioactivity associated with transfected cells or membrane from these cells. If the compound binds to the receptor, the binding of labeled ligand to the receptor is inhibited as determined by a reduction of labeled ligand which binds to the receptors. This method is called binding assay. Naturally, this same technique can be used to look for an agonist.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve measuring or, qualitatively or quantitatively, detecting the competition of binding of a candidate compound to the polypeptide with a labeled competitor (*e.g.* agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring human AXOR29 receptor activity in the mixture, and comparing the AXOR29 activity of the mixture to a control mixture which contains no candidate compound.

Another screening procedure involves the use of mammalian cells (CHO, HEK 293, *Xenopus* Oocytes, RBL-2H3, *etc.*) that are transfected to express human AXOR29 receptor. The cells are loaded with an indicator dye that produces a fluorescent signal when bound to calcium, and the cells are contacted with a test substance and a receptor agonist, such as S-1-P. Any change in

- 18 -

fluorescent signal is measured over a defined period of time using, for example, a fluorescence spectrophotometer or a fluorescence imaging plate reader. A change in the fluorescence signal pattern generated by the ligand indicates that a compound is a potential antagonist or agonist for the receptor.

5 Another screening procedure involves use of mammalian cells (CHO, HEK293, *Xenopus* Oocytes, RBL-2H3, *etc.*) that are transfected to express the receptor of interest, and that are also transfected with a reporter gene construct that is coupled to activation of the receptor (for example, luciferase or beta-galactosidase behind an appropriate promoter). The cells are contacted with a test substance and the receptor agonist (ligand), such as S-1-P or di-hydro S-1-P, and the signal
10 produced by the reporter gene is measured after a defined period of time. The signal can be measured using a luminometer, spectrophotometer, fluorimeter, or other such instrument appropriate for the specific reporter construct used. Inhibition of the signal generated by the ligand indicates that a compound is a potential antagonist for the receptor.

15 Another screening technique for antagonists or agonists involves introducing RNA encoding the AXOR29 polypeptide into *Xenopus* oocytes (or CHO, HEK 293, RBL-2H3, *etc.*) to transiently or stably express the receptor. The receptor oocytes are then contacted with the receptor ligand, such as S-1-P or di-hydro S-1-P, and a compound to be screened. Inhibition or activation of the receptor is then determined by detection of a signal, such as, cAMP, calcium, proton, or other ions.

20 Another method involves screening for AXOR29 polypeptide inhibitors by determining inhibition or stimulation of AXOR29 polypeptide-mediated cAMP and/or adenylate cyclase accumulation or diminution. Such a method involves transiently or stably transfecting a eukaryotic cell with AXOR29 polypeptide receptor to express the receptor on the cell surface. The cell is then exposed to potential antagonists in the presence of AXOR29 polypeptide ligand, such as S-1-P or di-hydro S-1-P. The changes in levels of cAMP is then measured over a defined period of time, for
25 example, by radio-immuno or protein binding assays (for example using Flashplates or a scintillation proximity assay). Changes in cAMP levels can also be determined by directly measuring the activity of the enzyme, adenylyl cyclase, in broken cell preparations. If the potential antagonist binds the receptor, and thus inhibits AXOR29 polypeptide-ligand binding, the levels of AXOR29 polypeptide-mediated cAMP, or adenylate cyclase activity, will be reduced or increased.

30 Another screening method for agonists and antagonists relies on the endogenous pheromone response pathway in the yeast, *Saccharomyces cerevisiae*. Heterothallic strains of yeast can exist in two mitotically stable haploid mating types, MATa and MATa. Each cell type secretes a small peptide hormone that binds to a G-protein coupled receptor on opposite mating-type cells which triggers a MAP kinase cascade leading to G1 arrest as a prelude to cell fusion. Genetic alteration of

certain genes in the pheromone response pathway can alter the normal response to pheromone, and heterologous expression and coupling of human G-protein coupled receptors and humanized G-protein subunits in yeast cells devoid of endogenous pheromone receptors can be linked to downstream signalling pathways and reporter genes (e.g., U.S. Patents 5,063,154; 5,482,835; 5,691,188). Such genetic alterations include, but are not limited to: (i) deletion of the STE2 or STE3 gene encoding the endogenous G-protein coupled pheromone receptors; (ii) deletion of the FAR1 gene encoding a protein that normally associates with cyclin-dependent kinases leading to cell cycle arrest; and (iii) construction of reporter genes fused to the FUS1 gene promoter (where FUS1 encodes a membrane-anchored glycoprotein required for cell fusion). Downstream reporter genes can permit either a positive growth selection (e.g., histidine prototrophy using the FUS1-HIS3 reporter), or a colorimetric, fluorimetric or spectrophotometric readout, depending on the specific reporter construct used (e.g., b-galactosidase induction using a FUS1-LacZ reporter).

The yeast cells can be further engineered to express and secrete small peptides from random peptide libraries, some of which can permit autocrine activation of heterologously expressed human (or mammalian) G-protein coupled receptors (Broach, *et al.*, *Nature* 384: 14-16, 1996; Manfredi, *et al.*, *Mol. Cell. Biol.* 16: 4700-4709, 1996). This provides a rapid direct growth selection (e.g., using the FUS1-HIS3 reporter) for surrogate peptide agonists that activate characterized or orphan receptors. Alternatively, yeast cells that functionally express human (or mammalian) G-protein coupled receptors linked to a reporter gene readout (e.g., FUS1-LacZ) can be used as a platform for high-throughput screening of known ligands, fractions of biological extracts and libraries of chemical compounds for either natural or surrogate ligands. Functional agonists of sufficient potency (whether natural or surrogate) can be used as screening tools in yeast cell-based assays for identifying G-protein coupled receptor antagonists. For example, agonists will promote growth of a cell with FUS-HIS3 reporter or give positive readout for a cell with FUS1-LacZ. However, a candidate compound that inhibits growth or negates the positive readout induced by an agonist is an antagonist. For this purpose, the yeast system offers advantages over mammalian expression systems due to its ease of utility and null receptor background (lack of endogenous G-protein coupled receptors), which often interferes with the ability to identify agonists or antagonists.

The present invention also provides a method for identifying new ligands not known to be capable of binding to a human AXOR29 receptor polypeptide. The screening assays described above for identifying agonists may be used to identify new ligands.

The present invention also contemplates agonists and antagonists obtained from the above described screening methods.

- 20 -

Examples of potential human AXOR29 polypeptide receptor antagonists include peptidomimetics, synthetic organic molecules, natural products, antibodies, *etc.*, that bind to the receptor but do not elicit a second messenger response, such that the activity of the receptor is prevented.

5 Potential antagonists also include proteins which are closely related to the ligand of the AXOR29 polypeptide receptor, *i.e.*, a fragment of the ligand, which have lost biological function, and when they bind to human AXOR29 polypeptide receptor, elicit no response.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, and ligands for human AXOR29 polypeptides, comprising:

10 (a) a human AXOR29 polypeptide, preferably that of SEQ ID NO:2; and further preferably comprises labeled or unlabeled S-1-P or di-hydro S-1-P;

(b) a recombinant cell expressing a human AXOR29 polypeptide, preferably that of SEQ ID NO:2; and further preferably comprises labeled or unlabeled S-1-P or di-hydro S-1-P; or

15 (c) a cell membrane expressing a human AXOR29 polypeptide; preferably that of SEQ ID NO: 2; and further preferably comprises labeled or unlabeled S-1-P or di-hydro S-1-P.

It will be appreciated that in any such kit, (a), (b), or (c) may comprise a substantial component.

As noted above, a potential antagonist is a small molecule that binds to the human AXOR29 polypeptide, making it inaccessible to ligands such that normal biological activity is prevented.

20 Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules.

Potential antagonists also include soluble forms of a human AXOR29 polypeptide receptor, *e.g.*, fragments of the receptor, which bind to the ligand and prevent the ligand from interacting with membrane bound human AXOR29 polypeptide.

25 The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or
30 inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

- 21 -


Constitutively active polypeptides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring human AXOR29 receptor activity in the mixture, and comparing the human AXOR29 receptor activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and AXOR29 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, *J. Mol. Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J. Biol. Chem.*, 270(16):9459-9471 (1995).

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well microtiter plates but also emerging methods such as the nanowell method described by Schullek, *et al.*, *Anal Biochem.*, 246: 20-29 (1997).


Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

"S-1-P (sphingosine-1-phosphate)" refers to the sphingolipid metabolite having the structure:



"Di-hydro S-1-P" (di-hydro sphingosine-1-phosphate)" refers to the sphingolipid metabolite having the structure:



- 22 -

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide" generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified RNA or DNA.

"Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is a mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells.

"Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a

polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, 1-12, in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol*, 182, 626-646, 1990, and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci*, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO:1..

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical.

A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more) PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

"Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over the length of the sequences being compared.

- 25 -

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole
5 length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two
10 polypeptide chains, on a residue by residue basis, taking into account not only exact correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an associated "score" from which the "% similarity" of the two sequences can then be determined.

15 Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J, *et al*, *Nucleic Acids Res*, 12, 387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the %
20 similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (*J Mol Biol*, 147,195-197, 1981, *Advances in Applied Mathematics*, 2, 482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison,
25 GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Needleman and Wunsch (*J Mol Biol*, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively.
30 Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F, *et al.*, *J Mol Biol*, 215, 403-410, 1990, Altschul S F, *et al.*, *Nucleic Acids Res.*, 25:389-3402, 1997, available from the

National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at www.ncbi.nlm.nih.gov) and FASTA (Pearson W R, *Methods in Enzymology*, 183, 63-99, 1990; Pearson W R and Lipman D J, *Proc Nat Acad Sci USA*, 85, 2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package).

5 Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, *Proc. Nat. Acad Sci. USA*, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

10 Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

15 "Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

30 Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between these terminal

-27-

positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \bullet I),$$

in which:

n_a is the number of nucleotide or amino acid differences,

x_a is the total number of nucleotides or amino acids in SEQ ID NO:1 or SEQ ID NO:2, respectively,

I is the Identity Index ,

\bullet is the symbol for the multiplication operator, and

in which any non-integer product of x_a and I is rounded down to the nearest integer prior to subtracting it from x_a .

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the two sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

"Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

- 28 -

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

Examples

Example 1: Mammalian Cell Expression

The receptors of the present invention are expressed in either human embryonic kidney 293 (HEK293) cells or adherent dhfr CHO cells. To maximize receptor expression, typically all 5' and 3' untranslated regions (UTRs) are removed from the receptor cDNA prior to insertion into a pCDN or pCDNA3 vector. The cells are transfected with individual receptor cDNAs by lipofectin and selected in the presence of 400 mg/ml G418. After 3 weeks of selection, individual clones are picked and expanded for further analysis. HEK293 or CHO cells transfected with the vector alone serve as negative controls. To isolate cell lines stably expressing the individual receptors, about 24 clones are typically selected and analyzed by Northern blot analysis. Receptor mRNAs are generally detectable in about 50% of the G418-resistant clones analyzed.

Example 2 Ligand bank for binding and functional assays

A bank of over 600 putative receptor ligands has been assembled for screening. The bank comprises: transmitters, hormones and chemokines known to act via a human seven transmembrane (7TM) receptor; naturally occurring compounds which may be putative agonists for a human 7TM receptor, non-mammalian, biologically active peptides for which a mammalian counterpart has not yet been identified; and compounds not found in nature, but which activate 7TM receptors with unknown natural ligands. This bank is used to initially screen the receptor for known ligands, using both functional (i.e. calcium, cAMP, microphysiometer, oocyte electrophysiology, etc, see below) as well as binding assays.

Example 3: Ligand Binding Assays

Ligand binding assays provide a direct method for ascertaining receptor pharmacology and are adaptable to a high throughput format. The purified ligand for a receptor is radiolabeled to high specific activity (50-2000 Ci/mmol) for binding studies. A determination is then made that the process of radiolabeling does not diminish the activity of the ligand towards its receptor. Assay conditions for buffers, ions, pH and other modulators such as nucleotides are optimized to establish a workable signal to noise ratio for both membrane and whole cell receptor sources. For these assays, specific receptor

- 29 -

binding is defined as total associated radioactivity minus the radioactivity measured in the presence of an excess of unlabeled competing ligand. Where possible, more than one competing ligand is used to define residual nonspecific binding.

Example 4: Functional Assay in *Xenopus* Oocytes

5 Capped RNA transcripts from linearized plasmid templates encoding the receptor cDNAs of the invention are synthesized in vitro with RNA polymerases in accordance with standard procedures. In vitro transcripts are suspended in water at a final concentration of 0.2 mg/ml. Ovarian lobes are removed from adult female toads, Stage V defolliculated oocytes are obtained, and RNA transcripts (10 ng/oocyte) are injected in a 50 nl bolus using a microinjection apparatus. Two electrode voltage clamps
10 are used to measure the currents from individual *Xenopus* oocytes in response to agonist exposure. Recordings are made in Ca²⁺ free Barth's medium at room temperature. The *Xenopus* system can be used to screen known ligands and tissue/cell extracts for activating ligands.

Example 5: Microphysiometric Assays

15 Activation of a wide variety of secondary messenger systems results in extrusion of small amounts of acid from a cell. The acid formed is largely as a result of the increased metabolic activity required to fuel the intracellular signaling process. The pH changes in the media surrounding the cell are very small but are detectable by the CYTOSENSOR microphysiometer (Molecular Devices Ltd., Menlo Park, CA). The CYTOSENSOR is thus capable of detecting the activation of a receptor which is coupled to an energy utilizing intracellular signaling pathway such as the G-protein coupled receptor
20 of the present invention.

Example 6: Extract/Cell Supernatant Screening

A large number of mammalian receptors exist for which there remains, as yet, no cognate activating ligand (agonist). Thus, active ligands for these receptors may not be included within the ligands banks as identified to date. Accordingly, the 7TM receptor of the invention is also functionally
25 screened (using calcium, cAMP, microphysiometer, oocyte electrophysiology, etc., functional screens) against tissue extracts to identify natural ligands. Extracts that produce positive functional responses can be sequentially subfractionated until an activating ligand is isolated identified.

Example 8: Calcium and cAMP Functional Assays

7TM receptors which are expressed in HEK 293 cells have been shown to be coupled
30 functionally to activation of PLC and calcium mobilization and/or cAMP stimulation or inhibition. Basal calcium levels in the HEK 293 cells in receptor-transfected or vector control cells were observed to be in the normal, 100 nM to 200 nM, range. HEK 293 cells expressing recombinant receptors are loaded with fura 2 and in a single day > 150 selected ligands or tissue/cell extracts are evaluated for

- 30 -

agonist induced calcium mobilization. Similarly, HEK 293 cells expressing recombinant receptors are evaluated for the stimulation or inhibition of cAMP production using standard cAMP quantitation assays. Agonists presenting a calcium transient or cAMP fluctuation are tested in vector control cells to determine if the response is unique to the transfected cells expressing receptor.

5 Example 9: Calcium Mobilization Assay for Human AXOR29

10 Based on the results generated in Figure 3, it was shown that an S-1-P exhibited an endogenous response in RBL 2H3 cells only at very high concentrations. Stable cell lines of the human AXOR29 receptor were prepared in the RBL 2H3 cell line. The expression of functionally active clones were followed using Fluorescent Imaging Plate Reader (FLIPR). The concentration-dependent responses for three clones to S-1-P is presented in Figure 3. As can be seen from this figure, the best clones respond in a concentration-dependent manner with EC₅₀s of about 120 nM. The best clones were characterized further, and that is shown in Figure 4. The cells responded with high potency through the human AXOR29 receptor to S-1-P. The cells also responded to dihydro-S-1-P. The cells did not respond to LPA, the other EDG family receptor ligand.

Claims

1. An isolated polypeptide selected from the group consisting of:

(a) an isolated polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;

5 (b) an isolated polypeptide comprising a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;

(c) an isolated polypeptide having at least 95% identity to the polypeptide sequence of SEQ ID NO:2; and

(d) fragments and variants of such polypeptides in (a) to (c).

10

2. The isolated polypeptide as claimed in Claim 1 comprising the polypeptide sequence of SEQ ID NO:2.

15 3. The isolated polypeptide as claimed in Claim 1 which is the polypeptide sequence of SEQ ID NO:2.

4. An isolated polynucleotide selected from the group consisting of:

(a) an isolated polynucleotide comprising a polynucleotide sequence having at least 95% identity to the polynucleotide sequence of SEQ ID NO:1;

20 (b) an isolated polynucleotide having at least 95% identity to the polynucleotide of SEQ ID NO:1;

(c) an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;

(d) an isolated polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;

25 (e) an isolated polynucleotide with a nucleotide sequence of at least 100 nucleotides obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof having at least 15 nucleotides;

(f) a polynucleotide which is the RNA equivalent of a polynucleotide of (a) to (e);

or a polynucleotide sequence complementary to said isolated polynucleotide

- 32 -

and polynucleotides that are variants and fragments of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.

5. An isolated polynucleotide as claimed in claim 4 selected from the group consisting of:

- 5 (a) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1;
- (b) the isolated polynucleotide of SEQ ID NO:1;
- (c) an isolated polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2; and
- (d) an isolated polynucleotide encoding the polypeptide of SEQ ID NO:2.

10

6. An expression system comprising a polynucleotide capable of producing a polypeptide of claim 1 when said expression vector is present in a compatible host cell.

15

7. A recombinant host cell comprising the expression vector of claim 6 or a membrane thereof expressing the polypeptide of claim 1.

8. A process for producing a polypeptide of claim 1 comprising the step of culturing a host cell as defined in claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.

20

9. An antibody immunospecific for the polypeptide of any one of claims 1 to 3.

10. A method for identifying agonist or antagonist of the polypeptide as claimed in Claim 2 comprising the steps of:

25

(a) contacting a cell expressing on the surface thereof the polypeptide, said polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and

- 33 -

(b) determining whether the compound binds to and activates or inhibits the polypeptide by measuring the level of a signal generated from the interaction of the compound with the polypeptide.

11. The method as claimed in Claim 10, wherein the ligand is labeled or unlabeled sphingosine-1-phosphate (S-1-P) or di-hydro sphingosine-1-phosphate (di-hydro-S-1-P).

12. A method for identifying an agonist or antagonist of the polypeptide as claimed Claim 2 comprising the steps of:

determining the inhibition of binding of a ligand to cells expressing the polypeptide on the surface thereof, or to cell membranes containing the polypeptide, in the presence of a candidate compound under conditions to permit binding to the polypeptide, and determining the amount of ligand bound to the polypeptide, such that a compound capable of causing reduction of binding of a ligand is an agonist or antagonist.

13. The method as claimed in Claim 12, wherein the ligand is labeled or unlabeled S-1-P or di-hydro S-1-P.

14. A method for screening to identify compounds that stimulate or that inhibit a function or level of the polypeptide as claimed in Claim 2, comprising a method selected from the group consisting of:

(a) measuring or, quantitatively or qualitatively, detecting the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;

(b) measuring the competition of the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor, preferably S-1-P or di-hydro S-1-P;

(c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;

- 34 -

(d) mixing a candidate compound with a solution comprising said polypeptide to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a to a control mixture which contains no candidate compound; or

(e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and said polypeptide in cells.

5

15. An antagonist identified by the method Claims 14.

16. An agonist identified by the method of Claim 14.

Figure 1

ATGGAGTCGGGGCTGCTGCGGCCGGCGCCGGTGAGCGAGGTTCATCGTCCTGCATTACAAC
TACACCGGCAAGCTCCGCGGTGCGCGCTACCAGCCGGGTGCCGGCCTGCGCGCCGACGCC
GTGGTGTGCCTGGCGGTGTGCGCCTTCATCGTGCTAGAGAACTAGCCGTGTGTTGGTG
CTCGGACGCCACCCGCGCTTCCACGCTCCCATGTTCTCTGCTCCTGGGCAGCCTCACGTTG
TCGGATCTGCTGGCAGGCGCCGCTACGCCGCCAACATCCTACTGTGCGGGCCGCTCACG
CTGAAACTGTCCCCGCGCTCTGTTTCGCACGGGAGGGAGGCGTCTTCGTGGCACTCACT
GCGTCCGTGCTGAGCCTCCTGGCCATCGCGCTGGAGCGCAGCCTCACCATGGCGCGCAGG
GGCCCCGCGCCGTCTCCAGTCGGGGGCGCACGCTGGCGATGGCAGCCGCGCGCTGGGGC
GTGTCGCTGCTCCTCGGGCTCCTGCCAGCGCTGGGCTGGAATTGCCTGGGTGCGCTGGAC
GCTTGCTCCACTGTCTTGCCGCTCTACGCCAAGGCCTACGTGCTCTTCTGCGTGCTCGCC
TTCGTGGGCATCCTGGCCGCTATCTGTGCACTCTACGCGCGCATCTACTGCCAGGTACGC
GCCAACGCGCGCGCCTGCCGCGACGGCCCGGACTGCGGGGACCACCTCGACCCGGGCG
CGTCGCAAGCCGCGCTCGCTGGCCTTGCTGCGCACGCTCAGCGTGGTGCTCCTGGCCTTT
GTGGCATGTTGGGGCCCCCTCTTCTGCTGCTGTTGCTCGACGTGGCGTGCCCGGCGCGC
ACCTGTCTGTACTCCTGCAGGCCGATCCCTTCTTGGGACTGGCCATGGCCAACTCACTT
CTGAACCCCATCATCTACACGCTCACCAACCGCGACCTGCGCCACGCGCTCCTGCGCCTG
GTCTGCTGCGGACGCCACTCCTGCGGCAGAGACCCGAGTGGCTCCAGCAGTCGGCGAGC
GCGGCTGAGGCTTCCGGGGGCTGCGCCGCTGCCTGCCCCCGGGCCTTGATGGGAGCTTC
AGCGGCTCGGAGCGCTCATCGCCCCAGCGCGACGGGCTGGACACCAGCGGCTCCACAGGC
AGCCCCGGTGACCCACAGCCGCCCGGACTCTGGTATCAGAACCGGCTGCAGACTGA

Figure 2

MESGLLRPAPVSEVIVLHYNVTGKLRGARYQPGAGLRADAVVCLAVCAFI VLENLAVLLV
LGRHPRFHAPMFLLLGSLTSLDLAGAAYAANILLSGPLTLKLSPALWFAREGGVFVALT
ASVLSLLAIALERSLTMARPGAPVSSRGRTLAMAAAAGVSLLLGLLPALGWNCLGRLD
ACSTVLPYAKAYVLFVLA FVGILAAICALYARIYCQVRANARRLPARPGTAGTTSTRA
RRKPRSLALLRTL SVVLLAFVACWGPLFLLLLLDVACPARTCPVLLQADPFLGLAMANS
LNPIIYTLTNRDLRHALLRLVCCGRHSCGRDPSSGSQASAAEASGGLRRCLPPGLDGSF
SGSERSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD

Figure 3





Figure 4



SEQUENCE LISTING

<110> SmithKline Beecham Corporation
SmithKline Beecham plc

<120> POLYNUCLEOTIDE AND POLYPEPTIDE SEQUENCES
OF HUMAN AXOR29 RECEPTOR AND METHODS OF SCREENING FOR
AGONISTS AND ANTAGONISTS OF THE INTERACTION BETWEEN HUMAN
AXOR29 RECEPTOR AND ITS LIGAND

<130> GP-50012WO

<140> TO BE ASSIGNED

<141> 2000-07-13

<150> 9916417.0

<151> 1999-07-13

<160> 2

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1197

<212> DNA

<213> HOMO SAPIENS

<400> 1

atggagtcgg	ggctgctg	gccggcgccg	gtgagcgagg	tcacgctcct	gcattacaac	60
tacaccggca	agctccgagg	tgcgcgctac	cagccgggtg	ccggcctg	cgccgacgcc	120
gtggtgtgcc	tggcggtgtg	cgccttcac	gtgctagaga	atctagccgt	gttggtggtg	180
ctcggacgcc	accgcgctt	ccacgctccc	atgttcctgc	tcctgggcag	cctcacgctg	240
tcggatctgc	tggcagggcg	cgcctacgcc	gccaacatcc	tactgtcg	gccgctcacg	300
ctgaaactgt	cccccgcgct	ctggttcgca	cgaggaggag	gcgtcttcgt	ggcactcact	360
gcgtccgtgc	tgagcctcct	ggccatcg	ctggagcgca	gcctcaccat	ggcgcgagg	420
gggcccgcgc	ccgtctccag	tcgggggcgc	acgtggcgca	tggcagccgc	ggcctggggc	480
gtgtcgctgc	tcctcgggct	cctgccagcg	ctgggctgga	attgcctggg	tcgcctggac	540
gcttgctcca	ctgtcttgcc	gctctacgcc	aaggcctacg	tgctcttctg	cgtgctcgcc	600
ttcgtgggca	tcctggccgc	tatctgtgca	ctctacgcgc	gcctctactg	ccaggtacgc	660
gccaacgcgc	ggcgctgcc	ggcacggccc	gggactgcgg	ggaccacctc	gacccggggc	720
cgctcgcaagc	cgcgctcgct	ggccttgctg	cgcacgctca	gcgtggtgct	cctggccttt	780
gtggcatggt	ggggccccct	cttctgctg	ctgttgctcg	acgtggcggtg	cccggcgcgc	840
acctgtcctg	tactcttgca	ggccgatccc	ttctggggac	tggccatggc	caactcactt	900
ctgaacccca	tcattctacac	gctcaccaac	cgcgacctgc	gccacgcgct	cctgcgcctg	960
gtctgctg	gacgccactc	ctgcggcaga	gacccgagtg	gctcccagca	gtcggcgagc	1020
gcggctgagg	cttcgggggg	cctgcgcgcg	tgctgtcccc	cgggccttga	tgggagcttc	1080
agcggctcgg	agcgtcatc	gccccagcgc	gacgggctgg	acaccagcgg	ctccacaggc	1140
agccccggtg	caccacacagc	cgccccggact	ctggtatcag	aaccggctgc	agactga	1197

<210> 2

<211> 398

<212> PRT

<213> HOMO SAPIENS

<400> 2

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1 5 10 15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
 20 25 30
 Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
 35 40 45
 Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60
 Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80
 Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser
 85 90 95
 Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
 100 105 110
 Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala
 115 120 125
 Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
 130 135 140
 Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly
 145 150 155 160
 Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
 165 170 175
 Gly Arg Leu Asp Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
 180 185 190
 Tyr Val Leu Phe Cys Val Leu Ala Phe Val Gly Ile Leu Ala Ala Ile
 195 200 205
 Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
 210 215 220
 Arg Leu Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
 225 230 235 240
 Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
 245 250 255
 Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
 260 265 270
 Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
 275 280 285
 Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
 290 295 300
 Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
 305 310 315 320
 Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
 325 330 335
 Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
 340 345 350
 Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
 355 360 365
 Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
 370 375 380
 Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
 385 390 395